

<https://helda.helsinki.fi>

---

## Dental age estimation in Somali children using the Willems et al. model

Metsäniitty, Mari

2018-11

---

Metsäniitty, M, Waltimo-Siren, J, Ranta, H, Fieuws, S & Thevissen, P 2018, 'Dental age estimation in Somali children using the Willems et al. model', International Journal of Legal Medicine, vol. 132, no. 6, pp. 1779-1786. <https://doi.org/10.1007/s00414-018-1926-1>

---

<http://hdl.handle.net/10138/260564>

<https://doi.org/10.1007/s00414-018-1926-1>

---

publishedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*



and Somalia (17%) were most often investigated. The first Somali asylum seekers arrived in Finland in the early 1990s following the mayhem of the civil war in Somalia. In 2016, there were 19,059 first- and second-generation Somalis in Finland [6].

The lack of African reference data, in particular from the Somali population, has been considered problematic in the legal age estimation context [7]. Although, scientifically, increasing evidence exists that ethnic differences are negligible in age estimations based on tooth development [8, 9], more confirmatory proof, especially related to comparisons including black populations, is needed. In fact, few studies of tooth development have been performed among sub-Saharan African populations [10–15]. Only Davidson and Rodd [6] have compared dental age and chronological age between Somali and Caucasian children.

Demirjian's dental age estimation method [16], constructed on a French-Canadian reference sample, is a widely used method to estimate age in children. After staging the development of the seven left mandibular permanent teeth (FDI 31–37), maturity scores are summed up and compared to reference tables or graphs. The age prediction performances of Demirjian's method have been validated in different populations and have resulted in the overestimation of chronological age [13, 18–20]. Willems et al. [18] adapted Demirjian's method using a large Belgian Caucasian reference sample and avoided calculating maturity scores. The Willems et al. model (WM) has been validated in multiple populations (Bangladeshi, Belgian Caucasian, Bosnian-Herzegovian, The Somali sample was used to validate the WM and to establish a Somali-specific age estimation model (SM) based on the WM. The SM was validated using leave-one-out cross-validation. Both models were analysed and compared on their

Table 1 Age and sex distribution of the Somali sample

Age/ years	Female (%)	Male (%)	Total
4–4.99	1 (0.32)	3 (0.93)	4
5–5.99	6 (1.93)	7 (2.16)	13
6–6.99	20 (6.43)	19 (5.86)	39
7–7.99	46 (14.79)	33 (10.19)	79
8–8.99	48 (15.43)	42 (12.96)	90
9–9.99	47 (15.11)	50 (15.43)	97
10–10.99	34 (10.93)	48 (14.81)	82
11–11.99	35 (11.25)	41 (12.65)	76
12–12.99	25 (8.04)	35 (10.80)	60
13–13.99	25 (8.04)	23 (7.10)	48
14–14.99	12 (3.86)	12 (3.70)	24
15–15.99	11 (3.54)	6 (1.85)	17
16–16.99	0 (0.00)	3 (0.93)	3
17–17.99	1 (0.32)	2 (0.62)	3
Total	311	324	635

## Material and methods

Ethical approval was granted by the Research Ethics Committee of the Hietalahti Institute, University of Helsinki (number 02/2010). All individuals eligible to be included in the current study were born in Finland after 1 January 1980. Both of an individual's parents had to be born in Somalia, their mother tongue had to be listed as Somali and their permanent address is Helsinki. According to the Finnish

**Table 2** Mean error (true age minus predicted age), mean absolute error and root mean squared error validating the Willems et al. model [48] in the collected Somali sample and the Somali model using leave-one-out cross-validation

Sex	N	Willems model			Somali model		
		ME (SD)	MAE (SD)	RMSE (95% CI)	ME (SD)	MAE (SD)	RMSE (95% CI)
F	307	0.20 (1.01)	0.78 (0.67)	1.02 (0.94;1.10)	0.04 (1.01)	0.77 (0.66)	1.01 (0.93;1.09)
M	321	−0.02 (1.00)	0.77 (0.63)	0.99 (0.92;1.07)	0.05 (1.04)	0.80 (0.67)	1.04 (0.96;1.12)
F + M	628	0.09 (1.01)	0.78 (0.65)	1.01 (0.95;1.06)	0.04 (1.03)	0.79 (0.66)	1.03 (0.97;1.08)

All reported ME, MAE and RMSE values are expressed in years. F, female; M, male; ME, mean error; SD, standard deviation; MAE, mean absolute error; RMSE, root mean square error and CI, confidence interval

age prediction performances, calculating the difference between true age and the predicted age. The mean error (ME) revealed statistically significant differences ( $p < 0.0001$ ) presented over- or underestimations (i.e., bias), the mean absolute error (MAE) differed 0.01 years in females ( $p < 0.05$ ) and 0.03 years in males ( $p < 0.05$ ) (Table 3). Figures 1, 2 and 3 present the quantified the magnitude of the errors (i.e., accuracy). Note that the MAE and RMSE reflect bias as well as lack of precision. The bias and the discrepancy between the age predictions of the WM and SM were evaluated using Wilcoxon signed rank tests. Spearman correlation between true age and the error in age estimation was used to evaluate if the direction and the magnitude of the difference depended on age: at younger ages, there was a tendency of overestimation, age. All statistical analyses were performed using SAS software, version 9.4 of the SAS System for Windows. (Spearman correlation between true age and the error in age estimation equalled 0.36,  $p < 0.0001$ ).

## Results

## Discussion

The intra-observer agreement was excellent, with Kappa and weighted Kappa values equal to 0.95 (95% confidence interval (CI) 0.92 to 0.98) and 0.98 (95% CI 0.96 to 0.99), respectively. The inter-observer agreement was also excellent, with Kappa and weighted Kappa values equal to 0.97 (95% CI 0.95 to 1.00) and 0.99 (95% CI 0.98 to 1.00), respectively.


The validated WM overestimated age for males with an ME of −0.02 years and underestimated age in the total sample (females + males) with an ME of 0.09 years, but these ME values were not significant ( $p = 0.27$  and  $p = 0.11$ , respectively). The WM significantly underestimated the age of females, with an ME of 0.20 years ( $p = 0.0006$ ). The MAE was 0.78 years in females, 0.77 years in males and 0.78 years in females and males combined. The RMSE was 1.02, 0.99, and 1.01 years for females, males and females and males combined, respectively (Table 2).

Validation of the SM resulted in a slight underestimation of age in the three respective sex groups, with MEs varying between 0.04 and 0.05 years, MAEs between 0.77 and 0.80 years and RMSEs between 1.01 and 1.04 years (Table 2).

**Table 3** Differences in mean error (true age minus predicted age), mean absolute error and root mean square error between the validated Willems et al. model and the Somali model

Sex	Difference bias		Difference accuracy		
	ΔME	p value	ΔMAE	ΔRMSE	p value
F	0.16	< 0.0001	0.01	0.01	0.97
M	−0.07	< 0.0001	−0.03	−0.05	0.0418
F + M	0.05	< 0.0059	−0.01	−0.02	0.18

All reported ME, MAE and RMSE values are expressed in years. F, female; M, male; ΔME, difference in mean error; ΔMAE, difference in mean absolute error; ΔRMSE, difference in root mean square error; P value P value from Wilcoxon signed rank test comparing the differences in true age and predicted age between the two models



performed locally. Phillips and van Wyk Kotze [14] constructed dental age-related tables in children of White and Coloured origin in South Africa (N = 1006), Indian (N = 234) and Black origin (N = 236; 171 Zulu and 65 Xhosa). The age range of the children under 16 years of age, resident in Sheffield, UK. Using Demirjian's method [17] to assess dental age, Somali children were significantly more advanced in dental development than the Caucasian children. The usefulness of the result is questionable, however, since it is unclear whether the exact age of Somali children was known, the Somali sample size was small.



Table 4 Root mean squared error calculated for females and males from the validated Willems et al. and the Somali models per age category of 1 year. Individuals with stage H were excluded

	Age/ years	N	RMSE		$\Delta$ RMSE
			Willems et al. model	Somali model	
F	4–4.99	1	−0.19 (−1.89, 1.51)	1.19 (−9.53, 11.91)	−1.38
	5–5.99	3	0.69 (0.20, 1.17)	0.67 (0.20, 1.55)	0.02
	6–6.99	19	0.45 (0.30, 0.60)	0.61 (0.40, 0.82)	−0.16
	7–7.99	46	0.66 (0.52, 0.80)	0.66 (0.52, 0.79)	0.00
	8–8.99	48	0.82 (0.65, 0.99)	0.82 (0.65, 0.99)	−0.00
	9–9.99	47	0.84 (0.66, 1.01)	1.08 (0.85, 1.30)	−0.24
	10–10.99	34	0.93 (0.70, 1.16)	1.00 (0.75, 1.24)	−0.07
	11–11.99	35	1.05 (0.80, 1.31)	1.12 (0.85, 1.39)	−0.07
	12–12.99	25	1.05 (0.74, 1.35)	0.95 (0.67, 1.22)	0.10
	13–13.99	25	1.29 (0.92, 1.67)	1.03 (0.73, 1.33)	0.26
	14–14.99	12	1.77 (0.98, 2.56)	1.35 (0.75, 1.95)	0.42
	15–15.99	11	1.95 (1.03, 2.86)	1.76 (0.93, 2.58)	0.19
	17–17.99	1	4.12 (32.88, 41.11)	3.99 (1.83, 39.80)	0.13
M	4–4.99	1	0.62 (−0.19, 1.42)	−0.14 (−1.41, 1.13)	0.76
	5–5.99	7	0.98 (0.36, 1.59)	1.09 (0.40, 1.77)	−0.11
	6–6.99	18	0.80 (0.53, 1.07)	0.79 (0.51, 1.06)	0.01
	7–7.99	33	0.80 (0.60, 1.00)	0.83 (0.62, 1.04)	−0.03
	8–8.99	42	0.72 (0.56, 0.88)	0.67 (0.52, 0.81)	0.05
	9–9.99	50	0.86 (0.69, 1.04)	0.90 (0.72, 1.08)	−0.04
	10–10.99	48	0.98 (0.78, 1.18)	1.02 (0.81, 1.23)	−0.04
	11–11.99	41	0.93 (0.73, 1.14)	0.93 (0.72, 1.14)	0.00
	12–12.99	35	1.24 (0.94, 1.55)	1.34 (1.02, 1.67)	−0.10
	13–13.99	23	1.03 (0.72, 1.35)	1.19 (0.83, 1.55)	−0.16
	14–14.99	12	0.88 (0.49, 1.27)	1.03 (0.57, 1.48)	−0.15
	15–15.99	6	1.78 (0.52, 3.04)	1.85 (0.54, 3.16)	−0.07
	16–16.99	3	2.12 (−0.63, 4.87)	2.24 (−0.68, 5.14)	−0.12
	17–17.99	2	2.86 (−3.29, 9.00)	3.00 (−3.45, 9.44)	−0.14

F, females; M, males; RMSE, root mean squared error;  $\Delta$ RMSE, difference in root mean square error. 95% confidence intervals for the RMSE are given between brackets

was reduced for this comparison from 635 (Table 2) to 628 (Table 2) subjects.

The good performance of validating the WM on the collected Somali children could partially be explained by the larger number of subjects in the Belgian Caucasian reference sample (WM,  $N=2116$ ; SM,  $N=635$ ) [18]. Larger samples tend to be a more accurate reflection of the population; hence, their sample means are more likely to be closer to the population mean. Even a small change in the number of subjects included in the considered sample can affect the outcome. In the present study, the validation sample was reduced from 311 to 307 subjects, increasing the ME of the WM from 0.20 to 0.21 years. The age prediction performances were not constant over age (Table 4). The higher the true age, the larger the absolute error was. The direction of the error depended on the true age: at younger ages, there was a tendency of overestimation and at older ages a tendency of underestimation (Spearman correlation between true age and the error in age estimation equalled 0.36,  $p<0.0001$ ). This is a classical finding in the application of regression models [34]. Since the present study was considering Somali children living in Finland and not in their home country, a possible influence of environmental factors might have an effect on dental maturity. Related to nutrition, Jääsaari et al. [34] longitudinally studied the association between dental maturity and body mass index (BMI) in an unselected group of Finnish children at the ages of 6 and 12 years. Tooth development was

registered according to Demirjian et al. [17] from the seven left mandibular permanent teeth (FDI-37) on 108 dental panoramic radiographs. Groups of delayed, average and advanced dental development were established and compared with normal and high BMI groups; energy intake was also studied with food records. They found no significant difference in BMI between delayed, normal or advanced dental maturity groups. However, the dental age in the advanced dental maturity group was advanced by 0.6 to 1.0 years when compared with the chronological age, and this group of children had a higher energy intake when compared with the average and delayed dental maturity groups ( $p < 0.004$ ). They conclude that there might be an association between advanced dental maturity and BMI. A previous study has also reported advanced dental development in overweight or obese children [32]. Converse results demonstrating a delay of dental development in underweight children have not, to the best of our knowledge, been published. Elamin and Liversidge [33] studied dental development in malnourished and normal BMI groups of young Arabs in Sudan. The subjects ( $n = 2115$ ) were between 2 and 22 years old. No significant difference in tooth formation was detected between the two groups. This study showed that teeth have substantial biological stability and are insulated from nutritional deprivation.

As a relative limitation of the study, the subjects included here had undergone dental panoramic radiography for valid clinical reasons related to dental health and deviations from normal occlusal development. Except for some longitudinal studies [31, 34, 35], this is typical for most, if not all, recent studies, making them comparable with each other. Poor dental health that leads to precocious extractions of primary teeth has been shown to speed up the eruption and alter the crown-to-root ratio of the succedaneous permanent teeth [37]. Nevertheless, dental developmental status is less affected than eruption status in subjects suffering from the premature loss of primary teeth [17]. Many of the children forming the material of the present study are likely to have exhibited malocclusion, since 95% of dental panoramic radiographs taken of children aged 7–12 years in the municipal healthcare centres in the City of Helsinki are taken for reasons related to orthodontics [38]. Although the study subjects were not analysed here for the presence or type of orthodontic problems, it is worth noting that individuals with discrepancies in jaw size have been shown to display advanced dental maturity in comparison to children with non-skeletal orthodontic problems; the difference reaches statistical significance in girls with mandibular prognathism [2].

## Conclusion

The small differences in age prediction performances of the WM and the constructed SM reflect the usefulness of the

Belgian population as a reference for forensic age predictions in Somali children living in Finland. The study also provides further support for the universal application of the WM to estimate age.

**Funding information** The Department of Oral and Maxillofacial Diseases, University of Helsinki, Finland funded the search of the sample from the Finnish Population Register Centre. The study was supported by a grant from the Association of Finnish Female Dentists. The authors are grateful to Senior Dental Officer Marja Noponen, DDS, PhD, and the personnel in the Unit for Specialised Oral Care in the Metropolitan Area and Kirkkonummi for arranging facilities for the study of radiographs and patient files. We thank Tuomo Maisala, DDS, in Oral Health Care, Helsinki, Finland for the search of the sample.

## Compliance with ethical standards

**Ethical approval** was granted by the Research Ethics Committee of the Hietel Institute, University of Helsinki (number 02/2010).

**Conflicts of interest** The authors declare that they have no conflicts of interest.

## References

- Schmeling A, Dettmeyer R, Rudolf E, Vieth V, Gesserick G (2016) Forensic age estimation: methods, certainty, and the law. *www.aerzteblatt-international.de/ref0416*
- Celikoglu M, Erdem A, Dane A, Demirci T (2011) Dental age assessment in orthodontic patients with and without skeletal malocclusions. *Orthod Craniofac Res* 14(2):52. <https://doi.org/10.1111/j.1601-6343.2011.01508.x>
- DuPlessis E, Araujo E, Behrents R, Kim K (2016) Relationship between body mass and dental and skeletal development in children and adolescents. *Am J Orthod Dentofac Orthop* 150(2):273. <https://doi.org/10.1016/j.ajodo.2015.12.031>
- Litsas G, Lucchese A (2016) Dental and chronological ages as determinants of peak growth period and its relationship with dental calcification stages. *Open Dent J* 10:998. <https://doi.org/10.2174/1874210601610010099>
- Metsäniitty M, Varkkola O, Waltimo-Sirén J, Ranta H (2017) Forensic age assessment of asylum seekers in Finland. *Int J Legal Med* 131:243250. <https://doi.org/10.1007/s00414-016-1498-x>
- Statistics Finland. Statistics Finland PX-Web databases. Population 31.12. by region, language, age, sex and type. [http://www.stat.fi/tup/suoluk/suoluk\\_vaesto\\_en.html](http://www.stat.fi/tup/suoluk/suoluk_vaesto_en.html)
- Gunawardena S, Liyanage U, Weeratna J, Mendis N, Perera H, Jayasekara R, Fernando R (2017) Forensic age estimation in anti-piracy trials in Seychelles: experiences and challenges faced. *Forensic Sci Int* 270:278.e278.e7. <https://doi.org/10.1016/j.forsciint.2016.11.010>
- Liversidge H, Smith B, Maber M (2010) Bias and accuracy of age estimation using developing teeth in 946 children. *Am J Phys Anthropol* 143:545554. <https://doi.org/10.1002/ajpa.21349>
- Thevissen P, Alqerban A, Asaumi J, Kahveci F, Kaur J, Kim YK, Pittaypat P, Van Vlierberghe M, Zhang Y, Fieuws S, Willems G (2010) Human dental age estimation using third molar developmental stages: accuracy of age predictions not using country specific information. *Forensic Sci Int* 201:10611
- Olze A, van Niekerk P, Schmidt S, Wernecke K-D, Rösing F, Gesserick G, Schmeling A (2006) Studies on the progress of third-



- molar mineralisation in a black African population. *Homo* 57:209–217
11. Phillips V, van Wyk Kotze T (2009) Dental age related tables for children of various ethnic groups in South Africa. *J Forensic Odontostomatol* 27:244
  12. Cavrč J, Galč I, Vodanović M, Brkić H, Gregov J, Viva S, Rey L, Cameriere R (2016) Third molar maturity index (I3M) for assessing age of majority in a black African population in Botswana. *Int J Legal Med* 130:1109–1120. <https://doi.org/10.1007/s00414-016-1344-1>
  13. Cavrč J, Vodanović M, Marušić A, Galč I (2016) Time of mineralization of permanent teeth in children and adolescents in Gaborone, Botswana. *Ann Anat* 203:22. <https://doi.org/10.1016/j.aanat.2015.08.001>
  14. Elamin F, Hector M, Liversidge H (2017) The timing of mandibular tooth formation in two African groups. *Ann Hum Biol* 44(3):261–272. <https://doi.org/10.1080/03014460.2016.1213313>
  15. Willems G, Lee S-S, Uys A, Bernitz H, Cadenas de Llano-Péruła M, Fieuws S, Thevissen P (2017) Age estimation based on Willems method versus new country-specific method in South African black children. *Int J Legal Med*. <https://doi.org/10.1007/s00414-017-1686-3>
  16. Davidson L, Rodd H (2001) Interrelationships between dental age and chronological age in Somali children. *Community Dent Health* 18:27–30
  17. Demirjian A, Goldstein H, Tanner J (1973) A new system of dental age assessment. *Hum Biol* 45:223–227
  18. Willems G, Van Olmen A, Spiessens B, Carels C (2001) Dental age estimation in Belgian children: Demirjian technique revisited. *J Forensic Sci* 46:893–895
  19. Maber M, Liversidge H, Hector M (2006) Accuracy of age estimation of radiographic methods using developing teeth. *Forensic Sci Int* 159S:S68–S73
  20. Flood S, Franklin D, Turlach B, McGeachie J (2013) A comparison of Demirjian's four dental development methods for forensic age estimation in South Australian sub-adults. *J Forensic Legal Med* 20:875–883. <https://doi.org/10.1016/j.jflm.2013.07.002>
  21. Galč I, Vodanović M, Cameriere R, Nakas E, Galč E, Selimović E, Brkić H (2011) Accuracy of Cameriere, Haavikko, and Willems radiographic methods on age estimation on Bosnian-Herzegovian children age groups 6–13. *Int J Legal Med* 125(2):323–325. <https://doi.org/10.1007/s00414-010-0515-8>
  22. Ye X, Jiang F, Sheng X, Huang H, Shen X (2014) Dental age assessment in 7–14-year-old Chinese children: comparison of Demirjian and Willems methods. *Forensic Sci Int* 244:436–438. <https://doi.org/10.1016/j.forsciint.2014.07.027>
  23. Ambarkova V, Galč I, Vodanović M, Biočina-Lukenda D, Brkić H (2014) Dental age estimations using Demirjian and Willems methods: cross sectional study on children from the Former Yugoslav Republic of Macedonia. *Forensic Sci Int* 234(187):e1–e7. <https://doi.org/10.1016/j.forsciint.2013.10.024>
  - Nik-Hussein N, Kee K, Gan P (2011) Validity of Demirjian and Willems methods for dental age estimation for Malaysian children aged 5–15 years old. *Forensic Sci Int* 204(1–2):e4–e6. <https://doi.org/10.1016/j.forsciint.2010.08.020>
  - Djukic K, Zelic K, Milenkovic P, Nedeljkovic N, Djuric M (2013) Dental age assessment validity of radiographic methods on Serbian children population. *Forensic Sci Int* 231:398–398.e5. <https://doi.org/10.1016/j.forsciint.2013.05.036>
  - Onat Altan H, Altan A, Bilgiç F, Akın Sözer Ö, Damlar (2016) The applicability of Willem's method for age estimation in southern Turkish children: a preliminary study. *J Forensic Legal Med* 38:24–27. <https://doi.org/10.1016/j.jflm.2015.11.015>
  - Akkaya N, Yilanci H, Göksülük D (2015) Applicability of Demirjian's four methods and Willems method for age estimation in a sample of Turkish children. *Legal Med* 17:355–359. <https://doi.org/10.1016/j.legalmed.2015.04.003>
  - Moorrees C, Fanning E, Hunt EE (1963) Age variation of formation stages for ten permanent teeth. *J Dent Res* 42:1490–1492
  - Thevissen PW, Fieuws S, Willems G (2010) Human third molars development: comparison of 9 country specific populations. *Forensic Sci Int* 201(1–3):102–105. <https://doi.org/10.1016/j.forsciint.2010.04.054>
  - Aykroyd R, Lucy D, Pollard A, Solheim T (1997) Technical note: regression analysis in adult age estimation. *Am J Phys Anthropol* 104(2):259–265
  - Jääsaari P, Tolvanen M, Niinikoski H, Karjalainen S (2016) Advanced dental maturity of Finnish 6 to 12-year-old children is associated with high energy intake. *Eur J Oral Sci* 124(5):465–471. <https://doi.org/10.1111/eos.12292>
  - Chehab DA, Tanbonliog T, Peyser J, Udin RT (2017) Association between body mass index and dental age in Hispanic children. *Gen Dent* 65(4):545–548
  - Elamin F, Liversidge H (2013) Malnutrition has no effect on the timing of human tooth formation. *PLoS One* 8(8):e72274. <https://doi.org/10.1371/journal.pone.0072274>
  - Nyström M, Kleemola-Kujala E, Evalahti M, Peck L, Kataja M (2001) Emergence of permanent teeth and dental age in a series of Finns. *Acta Odontol Scand* 59:456–458
  - Krekmanova L, Carlstedt-Duke J, Marcus C, Dahllöf G (1999) Dental maturity in children of short stature - a two-year longitudinal study of growth hormone substitution. *Acta Odontol Scand* 57(2):93–96
  - Brin I, Koyoumdjisky-Kaye E (1981) The influence of premature extractions of primary molars on the ultimate root length of their permanent successors. *J Dent Res* 60:962–965
  - Kim C, Hong Y, Han D, Hong H, Kim Y, Bae K (2011) A prospective cohort study on emergence of permanent teeth and caries experience in Korean children. *Int J Paediatr Dent* 21:260–269. <https://doi.org/10.1111/j.1365-263X.2011.01114.x>
  - Pakbaznejad Esmaeili E, Ekholm M, Haukka J, Waltimo-Sirén J (2016) Quality assessment of orthodontic radiography in children. *Eur J Orthod* 38:961–962